

MINISTRY OF EDUCATION AND SCIENCE UKRAINE
ODESSA NATIONAL I. I. MECHNIKOV UNIVERSITY

ФОТОЭЛЕКТРОНИКА

**PHOTOELECTRONICS
INTER-UNIVERSITIES SCIENTIFIC ARTICLES**

Founded in 1986

Number 22

Odessa
ONU
2013

« PHOTOELECTRONICS »
№ 22 – 2013

INTER-UNIVERSITIES SCIENTIFIC
ARTICLES

Founded in 1986

Certificate of State Registration
KB No 15953

« ФОТОЭЛЕКТРОНИКА »
№ 22–2013

МЕЖВЕДОМСТВЕННЫЙ НАУЧНЫЙ
СБОРНИК

Основан в 1986 г.

Свидетельство о Государственной
регистрации KB № 15953

The results of theoretical and experimental studies in problems of the semiconductor and microelectronic devices physics, opto- and quantum electronics, quantum optics, spectroscopy and photophysics of nucleus, atoms, molecules and solids are presented in the issue. New directions in the photoelectronics, stimulated by problems of the super intense laser radiation interaction with nuclei, atomic systems and substance, are considered.

For lecturers, scientists, post-graduates and students.

У збірнику наведено результати теоретичних і експериментальних досліджень з питань фізики напівпровідників та мікроелектронних приладів, опто- та квантової електроніки, квантової оптики, спектроскопії та фотофізики ядра, атомів, молекул та твердих тіл. Розглянуто нові напрямки розвитку фотоелектроніки, пов'язані із задачами взаємодії надінтенсивного лазерного випромінювання з ядром, атомними системами, речовиною.

Для викладачів, наукових працівників, аспірантів, студентів.

В сборнике приведены результаты теоретических и экспериментальных исследований по вопросам физики полупроводников и микроэлектронных приборов, опто- и квантовой электроники, квантовой оптики, спектроскопии и фотофизики ядра, атомов, молекул и твердых тел. Рассмотрены новые направления развития фотоэлектроники, связанные с задачами взаимодействия сверхинтенсивного лазерного излучения с ядром, атомными системами, веществом.

Для преподавателей, научных работников, аспирантов, студентов.

Editorial board «Photoelectronics»:

Editor-in-Chief **V. A. Smyntyna**

Kutalova M. I. (Odessa, Ukraine, responsible editor);

Vaxman Yu. F. (Odessa, Ukraine);

Litovchenko V. G. (Kiev, Ukraine);

Gulyaev Yu. V. (Moscow, Russia);

D'Amiko A. (Rome, Italy)

Mokrickiy V. A. (Odessa, Ukraine);

Neizvestny I. G. (Novosibirsk, Russia);

Starodub N. F. (Kiev, Ukraine);

Vikulin I. M. (Odessa, Ukraine)

The issue is introduced to the List of special editions of the Ukrainian Higher Certification Commission in physics-mathematics and technical sciences. Scientific articles «Photoelectronics» collection abstracted in ВИНІТИ and «Джерело», and are in scientificmetric base INDEX COPERNICUS with ICV 5.19.

Збірник включено до Переліку спеціальних видань ВАК України з фізико-математичних та технічних наук. Збірник «Photoelectronics» реферується у ВИНІТИ (Москва) та «Джерело» (Київ) і знаходиться у наукометричній базі INDEX COPERNICUS з ICV 5.19.

Сборник включен в Список специальных изданий ВАК Украины по физико-математическим и техническим наукам. Сборник «Photoelectronics» реферируется в ВИНІТИ (Москва) и «Джерело» (Киев) и находится в наукометричной базе INDEX COPERNICUS с ICV 5.19.

Address of editorial board:

Odessa National I. I. Mechnikov University 42, Pasteur str., Odessa, 65026, Ukraine

e-mail: wadz@mail.ru, tel.: +38-048765 32 16.

Information is on the site: <http://www.photoelectronics.onu.edu.ua>

TABLE OF CONTENTS:

<i>Valentyn A. Smyntyna</i>	5
<i>A. Tereshchenko, R. Viter, I. Konup, V. Ivanitsa , S. Geveliuk, Yu. Ishkov, V. Smyntyna</i> TiO ₂ -PORPHYRIN NANOSTRUCTURES FOR AMINO ACID DETECTION	8
<i>A. G. Kuzmich, D. A. Andrusenko, P. A. Tesel'ko, M. V. Isaiev, R. M. Burbelo</i> THERMOELASTIC STRESSES STUDY IN NANOCOMPOSITE SYSTEM “POROUS SILICON – LIQUID”	15
<i>A. V. Glushkov</i> OPERATOR PERTURBATION THEORY TO HYDROGEN ATOM IN A STRONG DC ELECTRIC FIELD	20
<i>O. Yu. Khetselius</i> NEW GEOMETRIC ATTRACTOR AND NEURAL NETWORKS APPROACH TO STUDYING CHAOTIC PROCESSES IN PHOTOELECTRONICS SYSTEMS	30
<i>O. O. Ptashchenko, F. O. Ptashchenko, V. R. Gilmudinova</i> TUNNEL SURFACE CURRENT IN GaAs P-N JUNCTIONS INDUCED BY AMMONIA MOLECULES ADSORPTION	38
<i>A. A. Svinarenko</i> SPECTROSCOPY OF AUTOIONIZATION STATES IN SPECTRA OF HELIUM, BARIUM AND LEAD ATOMS: NEW SPECTRAL DATA AND CHAOS EFFECT	43
<i>G. P. Prepelitsa</i> NON-LINEAR OPTICS AND SPECTROSCOPY OF ATOMIC AND LASER SYSTEMS WITH ELEMENTS OF A CHAOS	51
<i>V. V. Buyadzhi, S. V. Brusentseva, P. A. Zaichko</i> STUDYING ENSEMBLES OF INTERVALS OF THE PARKINSONIAN TREMOR AND LOCAL POTENTIAL FLUCTUATIONS ON THE BASIS OF THE THEORY OF CHAOS	61
<i>Iatsunskiy I. R., Smyntyna V. A., Pavlenko M. M.</i> DETECTION OF AMMONIA MOLECULES USING OPTICAL REFLECTANCE FROM NANOSTRUCTURED SILICON SURFACE	66
<i>A. P. Fedchuk, A. V. Glushkov, YA. I. Lepikh, L. Lovett, A. V. Ignatenko</i> STARK EFFECT AND RESONANCES IN THE IONIZATION CONTINUUM FOR EXCITONS IN QUANTUM DOTS AND ATOMS IN AN ELECTRIC FIELD	72

<i>Yu. F. Vaksman, Yu. A. Nitsuk, Yu. N. Purtov, A. S. Nasibov, P. V. Shapkin</i> OBTAINING AND OPTICAL PROPERTIES OF ZnS:Ti CRYSTALS	78
<i>N. Serga</i> RELATIVISTIC THEORY OF SPECTRA OF PIONIC ATOMS WITH ACCOUNT OF THE RADIATIVE AND NUCLEAR CORRECTIONS	84
<i>A. N. Shakhman</i> STRONG π NUCLEAT INTERACTION EFFECTS IN SPECTROSCOPY OF HADRONIC ATOMS	93
<i>V. A. Borschak., V. A. Smyntyna, IE. V. Brytavskiy, S. V. Zubritskiy, M. I. Kutalova, YA. I. Lepikh</i> MICROSTRUCTURAL FEATURES AND COMPONENTIAL ANALYSIS OF THIN FILM CdS-Cu ₂ S PHOTOSENSING STRUCTURES AS ELEMENT OF IMAGE SENSOR....	98
<i>G. P. Prepelitsa, V. V. Buyadzhi, V. B. Ternovsky</i> NON-LINEAR ANALYSIS OF CHAOTIC SELF-OSCILLATIONS IN BACKWARD- WAVE TUBE	103
<i>A. A. Kuznetsova, L. A. Vitavetskaya, YU. G. Chernyakova, D. A. Korchevsky</i> CALCULATING THE RADIATIVE VACUUM POLARIZATION CONTRIBUTION TO THE ENERGY SHIFT OF 2P-2S TRANSITION IN PIONIC DEUTERIUM.....	108
<i>V. A. Smyntyna, L. M. Filevska, O. V. Sviridova</i> TOPOLOGICAL FEATURES OF TIN DIOXIDE FILMS OBTAINED FROM THE BIS(ACETYLACETONATO)DICHLOROTIN COMPLEXES	112
<i>P. A. Kondratenko, Yu. M. Lopatkin, T. N. Sakun</i> ELECTRON STRUCTURE AND RELAXATION PROCESSES IN RESAZURIN IN A HIGHLY EXCITED STATE	117
<i>Yu. V. Dubrovskaya</i> ATOMIC CHEMICAL COMPOSITION EFFECT ON THE BETA DECAY PROBABILITIES.....	124
<i>Myndrul V. B., Bak A. YU., Karakis YU. N., Kutalova M. I., Zatovskaya N. P., Zubritskii S. V.</i> DETERMINATION IN MOBILITY OF NON-EQUILIBRIUM CARRIERS CONSIDERING MOTION VELOCITY DISTRIBUTION.....	130
<i>L. V. Nikola</i> AUGER-ELECTRON SPECTROSCOPY OF TRANSIENT METALS	135
<i>YU. M. Lopatkin, V. I. Mikhailenko, A. A. Svinarenko</i> SPECTROSCOPY OF THE SPECTRAL LINES BROADENING AND SHIFT FOR HEAVY ELEMENTS IN THE BUFFER GAS.....	139
<i>O. Yu. Khetselius</i> HYPERFINE STRUCTURE PARAMETERS OF THE HEAVY ISOTOPES: CONSISTENT NUCLEAR-QED THEORY	146
Інформація для авторів наукового збірника «Фотоелектроніка»	153

Kherson Department, Odessa National Polytechnical University, 1, Shevchenko pr., Ukraine
e-mail: quantsha@mail.ru

STRONG π NUCLEAT INTERACTION EFFECTS IN SPECTROSCOPY OF HADRONIC ATOMS

New New relativistic method on the basis of the Klein-Gordon-Fock equation with a generalized pion-nuclear potential is used to determine the strong interaction effects in spectroscopy of pionic atoms. The numerical data on 2p-1s, 3d-2p, 4f-3d, 5g-4f transition energies on the basis of the Klein-Gordon-Fock theory with the pion-nucleus potential, the Klein-Gordon-Fock theory with account of the radiative and the finite nuclear size corrections are presented for in some middle and heavy pionic atoms. It is performed comparison between theoretical data and results of measurements in the Berkley, CERN and Virginia laboratories.

1. Introduction

In this paper we goes on our studying on spectroscopy of pionic atoms and quantitative estimate of the strong pion-nucleon interactions contributions to the transitions energies in the X-ray spectra of pionic atomic systems. Let us note that the spectroscopy of hadron atoms has been used as a tool for the study of particles and fundamental properties for a long time. Exotic atoms are also interesting objects as they enable to probe aspects of atomic and nuclear structure that are quantitatively different from what can be studied in electronic or “normal” atoms. As it is known, one of the fundamental questions in the modern hadron’s physics is that the hadron masses being much higher than the mass of their quark content. The current mass of the up (u) and down (d) quarks is two orders of magnitude smaller than a typical hadron’s mass of about 1 GeV. This extraordinary phenomenon is proposed to originate from spontaneous breaking of chiral symmetry of massless quarks in strong interaction physics [1-5]. At present time one of the most sensitive tests for the chiral symmetry breaking scenario in the modern hadron’s physics is provided by studying the exotic hadron-atomic systems. Nowadays the transition energies in pionic (kaonic, muonic etc.) atoms are measured with an unprecedented precision and from studying spectra of the hadronic atoms it is possible to investigate the strong inter-

action at low energies measuring the energy and natural width of the ground level with a precision of few meV [1-10].

The mechanism of creation of the hadronic atoms is well known now (e.g. [1,2]). Really, such an atom is formed when a negative pion enters a medium, loses its kinetic energy through ionization and excitation of the atoms and molecules and eventually is captured, replacing the electron, in an excited atomic orbit. The further de-excitation scenario includes the different cascade processes such as the Auger transitions, Coulomb de-excitation, scattering etc. When a pion reaches a low-n state with the little angular momentum, strong interaction with the nucleus causes its absorption. The strong interaction is the reason for a shift in the energies of the low-lying levels from the purely electromagnetic values and the finite lifetime of the state corresponds to an increase in the observed level width. The possible energy shifts caused by the pion-induced fluorescence X-rays were checked in the measurement of the pion beams at PSI in Switzerland. For a long time the similar experimental investigations have been carried out in the laboratories of Berkley, Virginia (USA), CERN (Switzerland).

The most known theoretical models to treating the hadronic (pionic, kaonic, muonic, antiprotonic etc.) atomic systems are presented in refs. [1-5,7,8]. The most difficult aspects of the theo-

retical modeling are reduced to the correct description of pion-nuclear strong interaction [1-3] as the electromagnetic part of the problem is reasonably accounted for. Besides, quite new aspect is linked with the possible, obviously, very tiny electroweak and hyperfine interactions.

2. Relativistic approach to pionic atoms spectra

All available theoretical models to treating the hadronic (kaonic, pionic) atoms are naturally based on the using the Klein-Gordon-Fock equation [2,5], which can be written as follows :

$$m^2 c^2 \Psi(x) = \left\{ \frac{1}{c^2} [i\hbar \partial_t + eV_0(r)]^2 + \hbar^2 \nabla^2 \right\} \Psi(x) \quad (1)$$

where c is a speed of the light, \hbar is the Planck constant, and $\Psi_0(x)$ is the scalar wave function of the space-temporal coordinates. Usually one considers the central potential $[V_0(r), 0]$ approximation with the stationary solution:

$$\Psi(x) = \exp(-iEt/\hbar) \varphi(x), \quad (2)$$

where $\varphi(x)$ is the solution of the stationary equation:

$$\left\{ \frac{1}{c^2} [E + eV_0(r)]^2 + \hbar^2 \nabla^2 - m^2 c^2 \right\} \varphi(x) = 0 \quad (3)$$

Here E is the total energy of the system (sum of the mass energy mc^2 and binding energy e_0). In principle, the central potential V_0 naturally includes the central Coulomb potential, the vacuum-polarization potential, the strong interaction potential.

The most direct approach to treating the strong interaction is provided by the well known optical potential model (c.g. [2]). Practically in all papers the central potential V_0 is the sum of the following potentials. The nuclear potential for the spherically symmetric density $\rho(r|R)$ is [6,13]:

$$V_{nucl}(r|R) = -\left((1/r) \int_0^r dr' r'^2 \rho(r'|R) + \int_r^\infty dr' r' \rho(r'|R) \right) \quad (4)$$

The most popular Fermi-model approximation the charge distribution in the nucleus $\rho(r)$

(c.f.[12]) is as follows:

$$\tilde{n}(r) = \tilde{n}_0 \left[1 + \exp[(r-c)/a] \right], \quad (5)$$

where the parameter $a=0.523$ fm, the parameter c is chosen by such a way that it is true the following condition for average-squared radius:

$$\langle r^2 \rangle^{1/2} = (0.836 \times A^{1/3} + 0.5700) \text{fm.}$$

The effective algorithm for its definition is used in refs. [12] and reduced to solution of the following system of the differential equations:

$$V'_{nucl}(r, R) = \left(1/r^2 \int_0^r dr' r'^2 \rho(r', R) \right) \equiv \left(1/r^2 \right) y(r, R) \quad (6)$$

$$y'(r, R) = r^2 \rho(r, R), \quad (7)$$

$$\tilde{n}'(r) = (\tilde{n}_0/a) \exp[(r-c)/a] \left[1 + \exp[(r-c)/a] \right]^2 \quad (8)$$

with the corresponding boundary conditions. Another, probably, more consistent approach is in using the relativistic mean-field (RMF) model, which been designed as a renormalizable meson-field theory for nuclear matter and finite nuclei [13]. To take into account the radiation corrections, namely, the effect of the vacuum polarization we have used the generalized Ueling-Serber potential with modification to take into account the high-order radiative corrections [5,12]. The most difficult aspect is an adequate account for the strong interaction. In the pion-nucleon state interaction one should use the following pulse approximation expression for scattering amplitude of a pion on the "i" nucleon [2,3]:

$$f_i(r) = \left\{ b'_0 + b'_1(t\tau) + [c'_0 + c'_1(t\tau)] kk' \right\} \delta(r-r_i); \quad (9)$$

where t and τ are the isospines of pion and nucleon. The nucleon spin proportional terms of the kind $\sigma[kk']$ are omitted. The constants in (9) can be expressed through usual s-wave (α_{2T}) and p-wave ($\alpha_{2T,2J}$) scattering length (T and J -isospin and spin of the system πN). The corresponding parameters in the Compton wave length $\tilde{\lambda}_\pi$ terms are as follows:

$$b'_0 = (\alpha_1 + 2\alpha_3)/3 = -0.0017 \lambda_\pi.$$

$$b'_0 = (\alpha_3 - \alpha_1)/3 = -0.086 \lambda_\pi.$$

$$c'_0 = (4\alpha_{33} + 2\alpha_{13} + 2\alpha_{31} + \alpha_{11})/3 = -0.208 (\lambda_\pi)^3.$$

$$c'_1 = (2\alpha_{33} - 2\alpha_{13} + \alpha_{31} - \alpha_{11})/3 = -0.184 (\lambda_\pi)^3. \quad (10)$$

The scattering amplitude for pion on a nucleus can be further received as a coherent sum of the πN -scattering lengths. πN -scattering lengths. πN -scattering lengths. In approximation of the only s-wave interaction the corresponding potential can be written in the Dezer form:

$$V_N(r) = -2\pi h^2 \mu_\pi^{-1} [ZA^{-1}a_p + (A-Z)A^{-1}a_n] \rho(r). \quad (11)$$

The s-wave lengths of the $\pi^{-1}p$ -scattering $a_p = (2\alpha_1 + \alpha_3)/3$ и $\pi^{-1}n$ - scattering $a_n = \alpha_3$; scattering are introduced to Eq. (11). Because of the equality between $a_n = b'_0 + b'_1$ and $a_p = b'_0 - b'_1$ (with an opposite sign) the theoretical shift of

the s-level with $T = 0$ ($A = 2Z$) from Eq. (12) is much less than the observed shift. So, the more correct approximation must take into account the effects of the higher orders.

In whole the energy of the hadronic atom is represented as the sum:

$$E \approx E_{KG} + E_{FS} + E_{VP} + E_N; \quad (12)$$

Here E_{KG} -is the energy of a pion in a nucleus (Z, A) with the point-like charge (dominative contribution in (12)), E_{FS} is the contribution due to the nucleus finite size effect, E_{VP} is the radiation correction due to the vacuum-polarization effect, E_N is the energy shift due to the strong interaction V_N . The last contribution can be defined from the experimental energy values as:

$$E_N = E - (E_{KG} + E_{FS} + E_{VP}) \quad (13)$$

From the other side the strong pion-nucleus interaction contribution can be found from the solution of the Klein-Gordon equation with the corresponding meson-nucleon potential. In this case,

this contribution E_N is the function of the potentials (8)-(11) parameters.

3. Results and conclusions

In table 1 we present the data on the transition energies in some pionic atoms (from. Refs. [1-4]): the measured values from the Berkley, CERN and Virginia laboratories, the theoretical values for the $2p-1s$, $3d-2p$, $4f-3d$, $5g-4f$ pionic transitions ($E_{\#1}^N$ - values from the Klein-Gordon-Fock equation with the pion-nucleus potential [2]; E_{KGF} - values from the Klein-Gordon-Fock equation with account of radiative corrections (our data); E_{KS} - RMF finite nuclear size contribution (our data), $E_{\#2}^N$ - values from the Klein-Gordon-Fock equation with the generalized pion-nuclear potential [5] (our data).

Table 1. Transition energies (keV) in the spectra of some pionic atoms (see text)

Atom	E_{EXP} Berkley	E_{EXP} CERN	E_{KGF}	E_{FS}	$E_{\#1}^N$	$E_{\#2}^N$
Transition 4f-3d						
Ca ⁴⁰	72,352 ± 0.009		72,357	0,00	72,361	72,362
I ¹²⁷	519,1 ± 1,1	520,8 ± 0,8	516,50	-0,22	520,25	520,68
Cs ¹³³	560,5 ± 1,1	562,0 ± 1,5	556,80	-0,33	561,47	562,12
La ¹³⁹	603,6 ± 0,9	604,9 ± 2,0	597,88	-0,45	604,56	604,84
Transition 5g-4f						
Ta ¹⁸¹	453,1 ± 0,4	453,90 ± 0,20	450,67	-0,01	453,56	453,56
Au ¹⁹⁷	532,5 ± 0,5	533,16 ± 0,20	531,99	-0,025	532,87	532,87
Hg ²⁰¹	-	547,14 ± 0,25	545,730	-0,036	546,85	547,22
Bi ²⁰⁹	589,8 ± 0,9	590,06 ± 0,30	588,24	-0,05	589,75	590,16
Th ²³²	698,0 ± 0,6	698,4 ± 0,4	693,88	-0,17	697,20	698,25
U ²³⁸	731,4 ± 1,1	732,0 ± 0,4	725,78	-0,23	729,80	730,52

Note: E_{EXP} (Virginia): Ta¹⁸¹(5g-4f)=453.4±0.3; U²³⁸(5g-4f)=730.88±0.75;

It is easily to understand that when there is the close agreement between theoretical and experimental shifts, the corresponding energy levels are not significantly sensitive to strong pion-nuclear interaction, i.e the electromagnetic contribution is dominative. In the opposite situation the strong-interaction effect is very significant. The analysis of the presented data indicate on the necessity of the further more exact experimental investigations and further improvement of the pion-nuclear potential modelling. One can see that the contributions provided by the finite size effect should be accounted in a precise theory. Really, under availability of the "exact" values of the transitions energies one can perform the comparison of the theoretically and experimentally defined transition energies in the X-ray spectra in order to make a redefinition of the pion-nucleon model potential parameters using Eqs. (9)-(11). Taking into account the increasing accuracy of the X-ray pionic atom spectroscopy experiments, one can conclude that the such a way will make more clear the true values for parameters of the pion-nuclear potentials and correct the disadvantage of widely used parameterization of the potentials (9)-(11).

References

1. Deslattes R., Kessler E., Indelicato P., de Billy L., Lindroth E., Anton J., Exotic atoms // *Rev. Mod. Phys.* -2003.-Vol.75.-P.35-70.
2. Backenstoss G., Pionic atoms//*Ann.Rev. Nucl.Sci.*-1970.-Vol.20.-P.467-510.
3. Menshikov L I and Evseev M K, Some questions of physics of exotic atoms // *Phys. Uspekhi.*2001-Vol. 171.-P.150-184.
4. Deloff A., *Fundamentals in Hadronic Atom Theory.*- Singapore: World Sci., 2003.-352P.
5. Serga I. N., Dubrovskaya Yu. V., Kvasikova A. S., Shakhman A. N., Sukharev D. E., Spectroscopy of hadronic atoms: Energy shifts // *J.of Physics: C Ser.(IOP).*-2012.-Vol.397.-P.012013 (5p)
6. Scherer S., *Introduction to Chiral Perturbation Theory*//*Advances in Nuclear Physics*, Eds. J. W. Negele and E. W. Vogt (Berlin, Springer).-2003.-Vol.27.-P.5-50.
7. Schroder H., Badertscher A., Goudsmit P., Janousch M., Leisi H., Matsinos E., Sigg D., Zhao Z., Chatellard D., Egger J., Gabathuler K., Hauser P., Simons L., Rusi El Hassani A., The pion-nucleon scattering lengths from pionic hydrogen and deuterium//*Eur.Phys.J.*-2001.-Vol. C21.-P.473
8. Lyubovitskij V., Rusetsky A., π atom in ChPT: Strong energy-level shift// *Phys. Lett.B.*-2000.-Vol.494.-P.9-13.
9. Anagnostopoulos D., Biri S., Boissourdain V., Demeter M., Borchert G. et al-PSI, Low-energy X-ray standards from pionic atoms// *Nucl. Inst. Meth.B.*-2003.-Vol.205.-P.9-18.
10. Batty C. J., Eckhause M., Gall K. P., et al, Strong interaction effects in high Z- K^- atoms//*Phys. Rev. C.*-1989.-Vol.40.-P.2154-2160.
11. Kuznetsova A. A., Kvasikova A. S., Shakhman A. N., Vitavetskaya L. A., Calculating radiative vacuum polarization contribution to the energy shift of 2p-2s transition in m-hydrogen// *Photoelectronics* .- 2012.-N21.-P.66-67.
12. Glushkov A. V., Gauge-invariant QED perturbation theory approach to calculating nuclear electric quadrupole moments, hyperfine structure constants for heavy atoms and ions/ Glushkov A.V., Khetselius O.Yu., Sukharev D.E., Gurnitskaya E.P., Loboda A.V., Florko T.A., Lovett L.// *Frontiers in Quantum Systems in Chemistry and Physics* (Berlin, Springer).-2008.-Vol.18.- P.505-522.
13. Serot B. D., *Advances in Nuclear Physics* Vol. 16: *The Relativistic Nuclear Many Body Problem* / Serot B. D., Walecka J. D.-Plenum Press, New York, 1986.

The article is received in editorial 03.07.2013

UDC 539.182

A. N. Shakhman

STRONG p-NUCLEAT INTERACTION EFFECTS IN SPECTROSCOPY OF HADRONIC ATOMS

Abstract

New relativistic method on the basis of the Klein-Gordon-Fock equation with a generalized pion-nuclear potential is used to determine the strong interaction effects in spectroscopy of pionic atoms. The numerical data on 2p-1s, 3d-2p, 4f-3d, 5g-4f transition energies on the basis of the Klein-Gordon-Fock theory with the pion-nucleus potential and account of radiative and finite size nuclear corrections are presented for some middle and heavy pionic atoms. It is performed comparison between theoretical data and results of measurements in the Berkley, CERN and Virginia laboratories.

Key words: strong interaction, hadronic atom, relativistic theory

УДК 539.182

A. N. Шахман

ЭФФЕКТЫ СИЛЬНОГО ВЗАИМОДЕЙСТВИЯ В СПЕКТРОСКОПИИ АДРОННЫХ АТОМОВ

Резюме

Новый релятивистский метод на основе уравнения Клейна-Гордона-Фока с обобщенными пион-ядерным потенциалом применен для изучения эффектов сильного взаимодействия в спектроскопии пионных атомов. Представлены данные по энергиям 2p-1s, 3d-2p, 4f-3d, 5g-4f переходов для некоторых средних и тяжелых пионных атомов, полученные на основе уравнения Клейна-Гордона-Фока с пион-ядерным потенциалом с учетом радиационных и конечно-размерных ядерных поправок. Проведено сравнение теоретических данных с данными измерений в лабораториях Беркли, ЦЕРН и Вирджиния.

Ключевые слова: сильное взаимодействие, адронный атом, релятивистская теория

УДК 539.182

A. M. Шахман

ЕФЕКТИ СИЛЬНОЇ ВЗАЄМОДІЇ В СПЕКТРОСКОПІЇ АДРОННИХ АТОМІВ

Резюме

Новий релятивістський метод на основі рівняння Клейна-Гордона-Фока із узагальненим піон-ядерним потенціалом застосовано до опису ефектів сильної взаємодії в спектроскопії піонних атомів. Отримані чисельні дані про 2p-1s, 3d-2p, 4f-3d, 5g-4f енергії переходів для деяких середніх і важких піонних атомів, отримані на основі рівняння Клейна-Гордона-Фока з піон-ядерним потенціалом та урахуванням радіаційних та скінченно-розмірних ядерних поправок. Проведено порівняння теоретичних даних та результатів вимірювань в лабораторіях Берклі, ЦЕРН і Вірджинія.

Ключові слова: сильна взаємодія, адронний атом, релятивістська теорія